

Dissertation Summary

The Star Formation Rate density of the Universe at $z=0.24$ and $z=0.4$ from $H\alpha$

Sergio Pascual

Universidad Complutense de Madrid

Electronic mail: spr@astrax.fis.ucm.es

Thesis work conducted at: Universidad Complutense de Madrid

Ph.D. Thesis directed by: Jesús Gallego & Jaime Zamorano; Ph.D. Degree awarded:

2004-07-09

Received _____

Subject headings: Galaxies: distances and redshifts – Galaxies: evolution –
Galaxies: luminosity function, mass function

Knowledge of both the global star formation history of the Universe and the nature of individual star-forming galaxies at different look-back times are essential to our understanding of galaxy formation and evolution. Deep redshift surveys suggest star-formation activity increases by a order of magnitude from $z=0$ to $z \simeq 1$. To test directly whether substantial evolution in the star-formation activity has occurred we need to measure the star formation rate (SFR) density and the properties of the corresponding star-forming galaxy populations at different redshifts using similar techniques.

The main goal of this thesis work¹ is to extend the survey of emission-line galaxies of the *Universidad Complutense de Madrid* (UCM survey²) to higher redshifts.

The H α line (6563Å) luminosity, related to the number of massive stars, is a direct measurement of the current star formation rate. To compare directly with the H α -selected sample of the UCM survey, the selection of candidates is made through narrow-band filters. The study of the colors of different objects (stars and galaxies) in the system of narrow- and broad band filters shows strong increase of color as different emission lines enter the narrow band filter. It is thus possible to make a feasible selection of emission line galaxies using narrow band filters. The relations between the broad-band magnitude and narrow-broad color with the line flux and the equivalent width of the emission line are developed. These relations are used to obtain the luminosity of the emission line, assuming an average obscuration and a mean value for the doublet [N II] $\lambda\lambda$ 6548,6584.

The night-sky air-glow spectrum has substantial gaps around 8200Å and 9200Å, corresponding to H α redshifted respectively to $z \sim 0.24$ and $z \sim 0.4$. We have used two sets of filters. For the Wide Field Camera in Isaac Newton telescope at La Palma observatory, two narrow band filters (50Å width) and centered at those wavelength were designed. We also made use of a filter centered at 8200Å, 150Å width, with CAFOS, in the 2.2m telescope at Calar Alto observatory. Two samples of candidates at those redshifts have been built. The analysis of the objects is centered in the distribution of equivalent width and luminosities, when compared with the UCM sample. A important aspect of a survey with narrow-band filters is that it is open to contaminant galaxies with other emission lines (mainly [O III] $\lambda\lambda$ 4959,5007 and [O II] λ 3727) at higher redshifts. The fraction of

¹The original text is written in Spanish and is available at http://t-rex.fis.ucm.es/Publications/Repository/tesis_spr.ps.gz

²http://t-rex.fis.ucm.es/ucm_survey

contaminants is estimated analytically and found negligible at the relatively high line fluxes of the objects.

The SFR of the two samples is calculated assuming that every candidate selected in both filters (after stars removal) is a $H\alpha$ emitter. With this conditions, the SFRs are 3 times the SFR of the local Universe at $z \sim 0.24$ and 5 times at $z \sim 0.40$. A sub-sample selected with the 8200Å filter has been analyzed using photometric redshifts. We have found that some of the candidates appear at redshifts not covered with emission-lines, being misclassified objects. The luminosity density is then over-estimated in a factor ~ 1.2 . This correction is applied to the sample selected with the 9200Å filter and the SFR recalculated. The evolution of the SFR with redshift is found to be well fitted with a power-law $\rho \propto (1+z)^\beta$ with $\beta = 4.1 \pm 0.1$

Spectroscopic follow-up observations have been made for the same sub-sample. Of the objects with $H\alpha$ confirmed by photometric redshifts, 12 (60%) are spectroscopically confirmed. Two objects show the line $[O\ III]\lambda\lambda 4959,5007$ in emission, corresponding to a redshift about $z \sim 0.6$. The presence of these two objects, at the relatively bright line flux limit of the sub-sample, exhibit a stronger evolution in the luminosity functions than supposed in the analytical approach.

The galaxies confirmed with $H\alpha$ in emission are mainly starburst-like (9 objects) and only 3 objects are HII-like. Only one object is classified as a LINER. The masses, from rotation curves, are about the order of magnitude of the galaxies of the local Universe. These galaxies represent a population very similar to the local population of star-forming galaxies selected with $H\alpha$.